


contd.
B¹



(Ia)

[wherein:

ring C, R^1 , R^2 , n and Z are as defined hereinbefore with the provisos that R^2 is not hydrogen and that Z is not CH_2 or a direct bond; and

R^{2a} represents hydrogen, halogeno, C_{1-3} alkyl, trifluoromethyl, C_{1-3} alkoxy, C_{1-3} alkylsulphanyl, $-NR^{3a}R^{4a}$ (wherein R^{3a} and R^{4a} , which may be the same or different, each represents hydrogen or C_{1-3} alkyl), or $R^{5a}(CH_2)_{za}X^{1a}$ (wherein R^{5a} is a 4-, 5- or 6-membered saturated heterocyclic group with 1-2 heteroatoms, selected independently from O, S and N, which heterocyclic group may bear 1 or 2 substituents selected from oxo, hydroxy, halogeno, cyano, C_{1-4} cyanoalkyl, C_{1-4} alkyl, C_{1-4} hydroxyalkyl, C_{1-4} alkoxy, C_{1-4} alkoxy C_{1-4} alkyl, C_{1-4} alkylsulphonyl C_{1-4} alkyl, C_{1-4} alkoxycarbonyl, C_{1-4} aminoalkyl, C_{1-4} alkylamino, di(C_{1-4} alkyl)amino, C_{1-4} alkylamino C_{1-4} alkyl, di(C_{1-4} alkyl)amino C_{1-4} alkyl, C_{1-4} alkylamino C_{1-4} alkoxy, di(C_{1-4} alkyl)amino C_{1-4} alkoxy and a group $-(O-)_f(C_{1-4}alkyl)_g$ ringD (wherein f is 0 or 1, g is 0 or 1 and ring D is a 4-6-membered saturated heterocyclic group with 1-2 heteroatoms, selected independently from O, S and N, which cyclic group may bear one or more substituents selected from C_{1-4} alkyl), za is an integer from 0 to 4 and X^{1a} represents a direct bond, $-O-$, $-CH_2-$, $-S-$, $-SO-$, $-SO_2-$, $-NR^{6a}C(O)-$, $-C(O)NR^{7a}-$, $-SO_2NR^{8a}-$, $-NR^{9a}SO_2-$ or $-NR^{10a}-$ (wherein R^{6a} , R^{7a} , R^{8a} , R^{9a} and R^{10a} each independently represents hydrogen, C_{1-3} alkyl or C_{1-3} alkoxy C_{2-3} alkyl));

and salts thereof, and prodrugs thereof for example esters and amides, in the manufacture of a medicament for use in the production of an antiangiogenic and/or vascular permeability reducing effect in warm-blooded animals such as humans.